CONTENTS

(Abstracts/Contents Lists published in Analytical Abstracts, ASCA, Biochemical Abstracts, Biological Abstracts, Chemical Abstracts, Chemical Titles, Chromatography Abstracts, Current Contents/Physical, Chemical & Earth Sciences, Current Contents/Life Sciences, Deep-Sea Research/Part B: Oceanographic Literature Review, Excerpta Medica, Index Medicus, Mass Spectrometry Bulletin, PASCAL-CNRS, Referativnyi Zhurnal and Science Citation Index)

Publisher's Note 1

Interplay of hydrophobic and electrostatic interactions in biopolymer chromatography. Effects of salts on the retention of proteins
by W. R. Melander, Z. El Rassi and Cs. Horváth (New Haven, CT, U.S.A.) (Received February 1st, 1989) 3

Mathematical modelling of the peak in liquid chromatography
by J. Plicka, V. Svoboda, I. Kleinmann and A. Uhlířová (Prague, Czechoslovakia) (Received January 24th, 1989) 29

Optimization of the gas chromatographic separation of five-membered ring polyarenes with an admixed BPhBT liquid crystal–Dexsil 300 stationary phase
by G. M. Janini and N. T. Filfil (Kuwait, Kuwait) (Received January 17th, 1989) 43

Contribution of longitudinal diffusion to band broadening in liquid chromatography
by A. Berthod, F. Chartier and J.-L. Rocca (Villeurbanne, France) (Received January 20th, 1989) 53

Discussion of a controversial chiral recognition model
by W. H. Pirkle and J. E. McCune (Urbana, IL, U.S.A.) (Received February 1st, 1989) . 67

Resolution of a coeluting chromatographic pair using Kalman filtering
by T. Barker and S. D. Brown (Newark, DE, U.S.A.) (Received January 31st, 1989) . . . 77

Use of a centrifugal partition chromatography for assessing partition coefficients in various solvent systems
by N. El Tayar, A. Marston, A. Bechalany, K. Hostettmann and B. Testa (Lausanne, Switzerland) (Received January 20th, 1989) 91

π-Acceptor amide group for liquid chromatographic chiral separations with special emphasis on the 3,5-dinitrobenzoyl amide
by R. Däppen, H. R. Karfunkel and F. J. J. Leusen (Basle, Switzerland) (Received January 12th, 1989) 101

Ionization of DEAE-cellulose. Dependence of pK on ionic strength
by M. A. Smith and P. C. Gillespie (Provo, UT, U.S.A.) (Received January 5th, 1989) . . 111

Study of the lipophilic character of a series of β-carbolines
by G. L. Biagi (Bologna, Italy), M. C. Pietrogrande (Ferrara, Italy), A. M. Barbaro and M. C. Guerra (Bologna, Italy), P. A. Borea (Ferrara, Italy) and G. Cantelli Forti (Bologna, Italy) (Received January 31st, 1989) 121

Normalisation of high-performance liquid chromatography peak retention times for computerised comparison of wheat prolamin chromatograms
by H. C. Sapirstein, M. G. Scanlon and W. Bushuk (Winnipeg, Canada) (Received January 24th, 1989) 127

Analytical resolution of 4(5)-alkylated γ-(δ)-lactones by high-performance liquid chromatography on a silica-bonded chiral polyacrylamide sorbent. Chromatographic characterization of a stationary phase
by M. Huffer and P. Schreier (Würzburg, F.R.G.) (Received January 13th, 1989) 137

(Continued overleaf)

(*Contents continued*)

Gas chromatographic and sorption properties of macroporous methacrylate copolymers
by J. Hradil and F. Švec (Prague, Czechoslovakia), N. P. Platonova and L. D. Belyakova (Moscow, U.S.S.R.) and V. Maroušek (Prague, Czechoslovakia) (Received January 27th, 1989) 143

Use of crown ethers in gas chromatography
by Y. Jin and R. Fu (Beijing, China) and Z. Huang (Wuhan, China) (Received January 13th, 1989) 153

Analysis of solvent residues in pharmaceutical bulk drugs by wall-coated open tubular gas chromatography
by D. W. Foust and M. S. Bergren (Kalamazoo, MI, U.S.A.) (Received February 10th, 1989) 161

Separation of 4-(2-pyridylazo)resorcinolato metal chelates by micellar electrokinetic capillary chromatography
by T. Saitoh, H. Hoshino and T. Yotsuyanagi (Sendai, Japan) (Received January 30th, 1989) 175

Chromatographic approaches to the quality control of chiral proprionate anti-inflammatory drugs and herbicides
by B. Blessington, N. Crabb, S. Karkee and A. Northage (Bradford, U.K.) (Received February 14th, 1989) 183

Derivatization of N-methyl and cyclic amino acids with dimethylformamide dimethyl acetal
by M. F. Grubb and P. S. Callery (Baltimore, MD, U.S.A.) (Received January 20th, 1989) 191

Degradation and analysis of polyoxyethylene monoalkyl ethers in the presence of acetyl chloride and ferric chloride
by J. Szymanowski (Poznań, Poland) and P. Kusz, E. Dziwiński, H. Szewczyk and K. Pyżalski (Kedzierzyn-Koźle, Poland) (Received January 16th, 1989) 197

Mass spectral investigations on trichothecene mycotoxins. VII. Liquid chromatographic–thermospray mass spectrometric analysis of macrocyclic trichothecenes
by T. Krishnamurthy (Aberdeen Proving Ground, MD, U.S.A.), D. J. Beck and R. K. Isensee (Whitesboro, NY, U.S.A.) and B. B. Jarvis (College Park, MD, U.S.A.) (Received January 6th, 1989) 209

Sample-induced internal gradient of ionic strength in ion-exclusion microcolumn liquid chromatography
by K. Šlais (Brno, Czechoslovakia) (Received February 10th, 1989) 223

Formic acid as a milder alternative to trifluoroacetic acid and phosphoric acid in two-dimensional peptide mapping
by D. J. Poll and D. R. K. Harding (Palmerston North, New Zealand) (Received January 18th, 1989) 231

Determination of polycyclic aromatic hydrocarbons in lubricating oil base stocks using high-performance liquid chromatography and gas chromatography–mass spectrometry
by J.-P. F. Palmentier, A. J. Britten, G. M. Charbonneau and F. W. Karasek (Waterloo, Canada) (Received January 23rd, 1989) 241

Adsorption chromatography on cellulose. IV. Separation of D- and L-tryptophan and D- and L-methyltryptophan on cellulose with aqueous solvents
by A. O. Kuhn and M. Lederer (Lausanne, Switzerland) and M. Sinibaldi (Monterotondo Scalo, Italy) (Received November 15th, 1988) 253

Separation of *cis* and *trans* isomers of unsaturated fatty acids by high-performance liquid chromatography in the silver ion mode
by W. W. Christie and G. H. McG. Breckenridge (Ayr, U.K.) (Received November 10th, 1988) 261

Separation of major phospholipid classes by high-performance liquid chromatography and subsequent analysis of phospholipid-bound fatty acids using gas chromatography
by M. Seewald and H. M. Eichinger (Kranzberg, F.R.G.) (Received January 2nd, 1989) . 271

Analysis of warfarin and its metabolites by reversed-phase ion-pair liquid chromatography with fluorescence detection
by Y. W. J. Wong and P. J. Davis (Austin, TX, U.S.A.) (Received January 11th, 1989) . 281

Separation of fungal sterols by normal-phase high-performance liquid chromatography. Application to the evaluation of ergosterol biosynthesis inhibitors
by G. A. Peacock and M. W. Goosey (Wantage, U.K.) (Received January 24th, 1989) . . 293

Use of liquid chromatography in the synthesis of isoluminol-labelled medroxyprogesterone acetate and zeranol
by H. Koehler, L. Larocque and S. Sved (Ottawa, Canada) (Received February 3rd, 1989) 305

Determination of preservatives in cosmetic products. II. High-performance liquid chromatographic identification
by N. de Kruijf, A. Schouten, M. A. H. Rijk and L. A. Pranoto-Soetardhi (Zeist, The Netherlands) (Received January 18th, 1989) 317

Characterization of indirect photometry for the determination of inorganic anions in natural water by ion chromatography
by N. Chauret and J. Hubert (Montreal, Canada) (Received January 11th, 1989) 329

Isocratic elution of sodium, ammonium, potassium, magnesium and calcium ions by ion-exchange chromatography
by H. Sato (Yokohama-shi, Japan) (Received January 16th, 1989) 339

Capillary gas chromatographic–mass spectrometric determination of acid herbicides in soils and sediments
by T. Tsukioka (Nagano, Japan) and T. Murakami (Tokyo, Japan) (Received January 11th, 1989) 351

Rapid high-performance liquid chromatographic separation of barley malt α-amylase on cyclobond columns
by C. A. Henson and J. M. Stone (Madison, WI, U.S.A.) (Received January 24th, 1989) . 361

Quantitative analysis of thorium in plutonium using reversed-phase liquid chromatography and spectrophotometric detection
by V. T. Hamilton, W. D. Spall, B. F. Smith and E. J. Peterson (Los Alamos, NM, U.S.A.) (Received January 12th, 1989) 369

Notes

Influence of the cellulose content of cellulose gels on the electrophoretic mobility
by M. Wroński (Lódź, Poland) (Received January 24th, 1989) 378

Chromatographic studies of metal complexes. V. Thin-layer chromatography of some octahedral cobalt(III) and nickel(II) complexes
by R. K. Ray and M. K. Bandyopadhyay (Parganas, India) and G. B. Kauffman (Fresno, CA, U.S.A.) (Received January 21st, 1989) 383

Use of absorbance ratios in densitometric measurements for the characterization and identification of natural products of pharmacological interest
by M. I. Walash and O. M. Salama (Mansoura, Egypt) and M. M. Bishr (Saffat, Kuwait) (Received December 27th, 1989) 390

Simple method for the preparation of spherical agarose and composite gel particles
by E. Presecan, H. Porumb and I. Lascu (Cluj-Napoca, Romania) (Received January 31st, 1989) 396

(*Continued overleaf*)

(*Contents continued*)

Improvement of a carbon–nitrogen elemental analyser for marine samples collected on glass-fibre filters
by C. Bechemin, D. Delmas and M.-J. Garet (Nieul-sur-Mer, France) (Received January 30th, 1989) 399

Large-volume spotting apparatus and its application to the semi-quantitative determination of organochlorine pesticides in tea
by H. Wan (Hangzhou, China) (Received January 27th, 1989) 403

Traditional oriental medicines. I. Black Pearl: identification and chromatographic determination of some undeclared medicinal ingredients
by A. By, J. C. Ethier, G. Lauriault, M. LeBelle, B. A. Lodge, C. Savard, W.-W. Sy and W. L. Wilson (Ottawa, Canada) (Received February 9th, 1989) 406

Resolution of neuroactive non-protein amino acid enantiomers by high-performance liquid chromatography utilising pre-column derivatisation with *o*-phthaldialdehyde-chiral thiols. Application to 2-amino-ω-phosphonoalkanoic acid homologues and α-amino-β-N-methylaminopropanoic acid (β-methylaminoalanine)
by M. R. Euerby, L. Z. Partridge and P. B. Nunn (London, U.K.) (Received February 14th, 1989) 412

Application of correlation chromatography to the investigation of biopolymers in wines
by I. Sh. Shatirishvili (Tbilisi, U.S.S.R.) (Received January 16th, 1989) 420

High-performance liquid chromatography of brassinosteroids in plants with derivatization using 9-phenanthreneboronic acid
by K. Gamoh (Kochi-shi, Japan) and K. Omote, N. Okamoto and S. Takatsuto (Niigata, Japan) (Received January 24th, 1989) 424

Liquid chromatographic separation of racemates on acetylated or carbamoylated β-cyclodextrin-bonded stationary phases
by M. Tanaka and T. Shono (Osaka, Japan), D.-Q. Zhu (Dalian, China) and Y. Kawaguchi (Kanagawa, Japan) (Received February 15th, 1989) 429

Milligram-scale separation of optical isomers of 2-pentafluoroethylalanine and 2-trifluoromethylalanine by medium-performance reversed-phase chromatography
by J. W. Keller and K. Niwa (Fairbanks, AK, U.S.A.) (Received February 21st, 1989) . . 434

Isocratic reversed-phase high-performance liquid chromatographic assay for a cryptand–cryptate–free metal ion system
by W. A. Pettit, B. K. Swailes and R. E. Peterson (Iowa City, IA, U.S.A.) (Received February 27th, 1989) 440

Use of free-flow electrophoresis for the purification of components separated by ion-pair chromatography
by R. Kessler, H. J. Manz and G. Székely (Basle, Switzerland) (Received February 2nd, 1989) 444

Purification of yeast killer toxin KT28 by receptor-mediated affinity chromatography
by M. Schmitt and F. Radler (Mainz, F.R.G.) (Received January 27th, 1989) 448

Author Index 453

In articles with more than one author, the name of the author to whom correspondence should be addressed is indicated in the article heading by a 6-pointed asterisk (*)